

# A NEW SPECIES OF *DALMATINELLA* RADOMAN, 1973 (CAENOGASTROPODA: HYDROBIIDAE) FROM CROATIA

LUBOŠ BERAN<sup>1</sup>, ALEKSANDRA RYSIEWSKA<sup>2</sup>, SEBASTIAN HOFMAN<sup>3</sup>, ARTUR OSIKOWSKI<sup>4</sup>, ANDRZEJ FALNIOWSKI<sup>2</sup>

<sup>1</sup>Nature Conservation Agency of the Czech Republic, Regional Office Kokořínsko – Máchův kraj Protected Landscape Area Administration, Česká 149, CZ – 276 01 Mělník, Czech Republic

<sup>2</sup>Department of Malacology, Institute of Zoology and Biomedical Research, Jagiellonian University, Gronostajowa 9, 30–387 Kraków, Poland

<sup>3</sup>Department of Comparative Anatomy, Institute of Zoology and Biomedical Research, Jagiellonian University, Gronostajowa 9, 30–387 Kraków, Poland

<sup>4</sup>Department of Animal Reproduction, Anatomy and Genomics, University of Agriculture in Krakow, Mickiewicza 24/28, 30–059 Kraków, Poland.

**Abstract** A new species of *Dalmatinella Radoman, 1973* from Croatia, found at three localities: Donja Rošca and Trnbusi at Cetina River and Ploče, Baćina lakes (type locality), is described. The shell, female reproductive organs and penis are described and illustrated. New species is compared with the type species of *Dalmatinella*: *D. fluviatilis Radoman, 1973*, from its type locality: Jankovića Buk, waterfalls, the Zrmanja River, Croatia. The new species is characterised by somewhat different habitus of the penis, and by the shell with higher spire and aperture. The differences are confirmed by the principal component analysis (PCA). Mitochondrial cytochrome oxidase subunit I (COI) confirmed species distinctness of the new species (*p*-distance 0.039).

**Key words** Gastropod, taxonomy, reproductive organs, shell biometry, PCA, mtDNA, cytochrome oxidase, Balkans

## INTRODUCTION

*Dalmatinella fluviatilis* Radoman, 1973, the only representative of the genus *Dalmatinella* Radoman, 1973 described so far, can probably be considered one of the endemites of the Zrmanja River, in northern Croatia (Falniowski & Szarowska, 2013). Apart from this type locality (Fig. 1), it was reported from the lowest part of the Neretva River, between Kula and Opuzen (Radoman, 1983). Considering more than 200km distance between Zrmanja and Neretva, without any records along this distance, Falniowski & Szarowska (2013) raised doubt that the same species of *Dalmatinella* could be present at both localities *Dalmatinella* could be represented by the same species. Recently, one of us (LB) collected specimens of *Dalmatinella* at three new localities, unreported in Radoman (1983). The present paper deals with the systematic status of these gastropods.

## MATERIAL AND METHODS

Snails were collected at four localities (Fig. 1: localities' numbers as in the map):

1. Donja Rošca, 43°33'44,9"N, 16°43'03,8"E, the Cetina River in the Prančevići dam reser-

voir, stones on the bank of the dam reservoir (Fig. 2A), 275m a.s.l., 29.8.2018, about 40 specimens;

2. Trnbusi, 43°29'22,6"N, 16°48'59,8"E, the Cetina River to the east of hill Gradina (308m a.s.l.), stones on the bank of the river (Fig. 2B), 200m a.s.l., 22.8.2019, 1 specimen;

3. Ploče, 43°04'11,8"N, 17°25'22,2"E, the S part of lake Sladinac, one stone on the bank, Baćina lakes, 9m a.s.l., 22.8.2019, about 30 specimens; type locality;


4. Jankovića Buk waterfalls, 44°12'09,8"N, 15°43'16,9"E, Zaton Obrovački, the Zrmanja River (Figs 2C–D), 9m a.s.l. 4.6.2011, about 25 specimens, the locality and materials the same as in Falniowski & Szarowska (2013).

Snails were collected with a 0.5mm sieve and fixed in 80% analytically pure ethanol, replaced two times. Next, the snails were put in fresh 80% analytically pure ethanol and kept in -20°C temperature in a refrigerator.


The shells were photographed with a Canon EOS 50D digital camera, under a Nikon SMZ18 microscope. The dissections were done under a Nikon SMZ18 microscope with dark field, equipped with Nikon DS-5 digital camera, whose captured images were used to draw anatomical structures with a graphic tablet.



**Figure 1** Geographic distribution of the localities of *Dalmatinella*: *D. simonae* – red dots (3 – type locality); *D. fluviatilis* – black dot (type locality); localities' numbers – see the text.



**Figure 2** Localities of *Dalmatinella*: A–B – Cetina River: A – Prančevići dam reservoir, B – Trnbusi; C – Jankovića Buk waterfalls; D – Zrmanja River directly over the waterfalls, *Dalmatinella* was collected from this place.



**Figure 3** Morphometric measurements used in PCA analysis.

Seven morphometric parameters of the shell (Szarowska, 2006; Falniowski *et al.*, 2007) were measured (Fig. 3) by one person using a Nikon DS-5 digital camera and ImageJ image analysis software (Rueden *et al.*, 2017). The linear measurements were then logarithmically transformed; for angular measurements the arcsine transformation was applied. Principal component analysis (PCA), based on the matrix of correlation, was computed, applying a descriptive, non-stochastic approach. The original observations were projected into PC space, to show relationships between the specimens, without any classification given *a priori* (Falniowski, 2003; Rohlf, 1998). The transformations and PCA calculations were made by the ClustVis 2.0 (<https://biit.cs.ut.ee/clustvis/>) (Metsalu & Jaak, 2015). Shell character states after Hershler & Ponder (1998).

DNA was extracted from whole specimens; tissues were hydrated in TE buffer (3×10 min); then total genomic DNA was extracted with the SHERLOCK extraction kit (A&A Biotechnology), and the final product was dissolved in 20 µl of tris-EDTA (TE) buffer. The extracted DNA was stored at -80°C at the Department of Malacology, Institute of Zoology and Biomedical Research, Jagiellonian University in Kraków (Poland).

A fragment of mitochondrial cytochrome oxidase subunit I (COI) was sequenced. Details of PCR conditions, primers used and sequencing were given in Szarowska *et al.* (2016). Sequences were initially aligned in the MUSCLE (Edgar, 2004) Programme in MEGA 6 (Tamura *et al.*, 2013) and then checked in BioEdit 7.1.3.0 (Hall, 1999). Uncorrected p-distances were calculated in MEGA 6. The estimation of the proportion of invariant sites and the saturation test for entire data sets (Xia, 2000; Xia *et al.*, 2003) were performed using DAMBE (Xia, 2013). In the phylogenetic analysis, additional sequences from GenBank were used as reference (Table 1). The data were analysed using approaches based on Bayesian Inference (BI) and Maximum Likelihood (ML). We applied the GTR model whose parameters were estimated by RaxML (Stamatakis, 2014). In the BI analysis, the GTR +I +Γ model of nucleotide substitution was applied. Model was selected using MrModelTest 2.3 (Nylander, 2004). The Bayesian analyses were run using MrBayes v. 3.2.3 (Ronquist *et al.*, 2012) with default of most priors. Two simultaneous analyses were performed, each with 10,000,000 generations, with one cold chain and three heated chains, starting from random trees and sampling the trees every 1,000 generations. The first 25% of the trees were discarded as burn-in. The analyses were summarised as a 50% majority-rule tree. Convergence was checked in Tracer v. 1.5 (Rambaut & Drummond, 2009). The Maximum Likelihood analysis was conducted in RAxML v. 8.2.12 (Stamatakis, 2014) using the 'RAxML-HPC v.8 on XSEDE (8.2.12)' tool via the CIPRES Science Gateway (Miller *et al.*, 2010). We applied the GTR model which is the only nucleotide substitution model implemented in RaxML, whose parameters were estimated by RaxML (Stamatakis, 2014).

**Table 1** Taxa used for phylogenetic analyses with their GenBank accession numbers and references.

Species	COI GB numbers	References
<i>Agrafia wiktoriae</i> Szarowska et Falniowski, 2011	JF906762	Szarowska & Falniowski, 2011
<i>Alzoniella finalina</i> Giusti & Bodon, 1984	AF367650	Wilke <i>et al.</i> , 2001
<i>Anagastina zetavalis</i> (Radoman, 1973)	EF070616	Szarowska, 2006
<i>Avenionia brevis berenguieri</i> (Draparnaud, 1805)	AF367638	Wilke <i>et al.</i> , 2001
<i>Belgrandiella kuesteri</i> (Boeters, 1970)	MG551325	Osikowski <i>et al.</i> , 2018
<i>Ecrobia maritima</i> (Milaschewitsch, 1916)	KX355835	Osikowski <i>et al.</i> , 2016
<i>Daphniola louisi</i> Falniowski & Szarowska, 2000	KM887915	Szarowska <i>et al.</i> , 2014a
<i>Dalmatinella fluviatilis</i> Radoman, 1973	KC344541-42	Falniowski & Szarowska, 2013
<i>Fissuria boui</i> Boeters, 1981	AF367654	Wilke <i>et al.</i> , 2001
<i>Graecoarganiella parnassiana</i> Falniowski & Szarowska, 2011	JN202352	Falniowski & Szarowska, 2011b
<i>Graziana alpestris</i> (Frauenfeld, 1863)	AF367641	Wilke <i>et al.</i> , 2001
<i>Grossuana angeltsekovi</i> Glöer & Georgiev, 2009	KU201090	Falniowski <i>et al.</i> , 2016
<i>Hauffenia tellinii</i> (Pollonera, 1898)	KY087861	Rysiewska <i>et al.</i> , 2017
<i>Horatia klecakiana</i> Bourguignat 1887	KJ159128	Szarowska & Falniowski, 2014
<i>Islamia zermanica</i> (Radoman, 1973)	KU662362	Beran <i>et al.</i> , 2016
<i>Montenegrospeum bogici</i> (Pešić & Glöer, 2012)	KM875510	Falniowski <i>et al.</i> 2014
<i>Pontobelgrandiella</i> sp. Radoman, 1978	KU497024	Rysiewska <i>et al.</i> , 2016
<i>Pseudorientalia</i> Radoman, 1973	KJ920477	Szarowska <i>et al.</i> , 2014b
<i>Radomaniola curta</i> (Küster, 1853)	KC011814	Falniowski <i>et al.</i> , 2012a
<i>Sadleriana fluminensis</i> (Küster, 1853)	KF193067	Szarowska & Falniowski 2013
<i>Tanousia zrmanjiae</i> (Brusina, 1866)	KU041812	Beran <i>et al.</i> , 2015

## SYSTEMATICS

Family Hydrobiidae Stimpson, 1865

Subfamily Sadlerianinae Szarowska, 2006

Genus *Dalmatinella* Radoman, 1973

*Dalmatinella simonae* Beran et Rysiewska sp. n.

urn:lsid:zoobank.org:act:853CA97A-78CC-46E9-998C-48DB998E7AFB

**Holotype** Ethanol-fixed specimen (Fig. 3), collected at Ploče, 43°04'11,8"N, 17°25'22,2"E, the S part of lake Sladinac, one stone on the bank, Baćina lakes (Fig. 1 – locality 1), Museum of Natural History of the University of Wrocław, Poland, voucher number MNHW-1364.


**Paratypes** 20 ethanol-fixed specimens, collected at Ploče, 43°04'11.8"N, 17°25'22.2"E, the S part of Lake Sladinac, one stone on the bank, Baćina lakes, voucher number MNHW-1364.

**Diagnosis** Shell small, ovate-conic or conic, distinguished from similarly high and broad *D. fluviatilis* by proportionally higher spire, aperture higher and slightly broader, body whorl lower, and by the penis with smaller outgrowths.

**Description** *Shell* (Figs 3–4) ovate-conic or conic, up to 2.15mm tall, 4–5 whorls growing regularly, spire height approximately 86% width of shell. Holotype measurements: Table 2. Teleoconch whorls highly convex, evenly rounded. Aperture broad, ovate, very weakly angled ad apically, peristome continuous, in its upper part adjacent to body whorl. Parietal lip complete, adnate, umbilicus small and partly covered by lip. Outer lip fluted, orthocline. Shell glossy with very slightly marked growth lines, thin and translucent, periostracum whitish or pinkish.

**Measurements** of holotype and shells of the sequenced specimens: Table 2. Shell variability slight. Principal component analysis (PCA) of the measurements given in Table 2 (Fig. 5), of the shells presented in Fig. 4, illustrates variability and morphological distinctness of the two species of *Dalmatinella*, at all the four localities.


**Soft parts morphology and anatomy** Body yellowish with rather strong black pigmentation. Female reproductive organs (Fig. 6) with long and weakly swollen loop of oviduct, moderately large, spherical bursa copulatrix and two receptacula seminis, proximal (rs<sub>2</sub>) big and distal (rs<sub>1</sub>)



**Figure 4** Shell variability of *Dalmatinella*: A–G – *D. simonae* n. sp. (A – holotype; B–G – specimens whose haplotypes are the same as in Fig. 8: site 1: B – 2E23, C – 2E24, D – 2E22; site 2: E – 1Z25, F – 1Z26; site 3: G – 2E19); H–L – *Dalmatinella fluviatilis* Radoman, 1973 (after Falniowski & Szarowska, 2013). Bar equals 1mm.


**Table 2** Measurements of the shell. Measured parameters as shown in Fig. 3. Holotype and 2E19 from site 3, 2E22-24 – site 1; 1Z25-26 – site 2; M – mean, SD – standard deviation

	a	b	c	d	e	$\alpha$	$\beta$
<i>Dalmatinella simonae</i> ; n=7							
Holotype	1.94	0.96	0.84	0.72	0.79	97	16
2E23	1.90	0.91	0.76	0.74	0.72	97	17
2E24	1.74	0.87	0.74	0.60	0.72	97	17
2E22	2.15	0.97	0.82	0.94	0.72	89	16
1Z25	2.10	0.95	0.81	0.91	0.68	95	15
1Z26	1.99	0.89	0.85	0.86	0.72	95	16
2E19	1.83	0.88	0.74	0.77	0.67	93	19
M	1.95	0.92	0.79	0.79	0.72	94.71	16.57
SD	0.144	0.041	0.047	0.120	0.039	2.928	1.272
Min	1.74	0.87	0.74	0.60	0.67	89	15
Max	2.15	0.97	0.85	0.94	0.79	97	19
<i>Dalmatinella fluviatilis</i> ; n=6							
D. f. 1	2.21	0.95	0.74	1.05	0.66	96	20
D. f. 2	2.09	0.93	0.73	0.97	0.60	96	14
D. f. 3	1.93	0.90	0.70	0.87	0.59	96	16
D. f. 4	1.83	0.88	0.71	0.87	0.58	96	18
D. f. 5	1.98	0.86	0.71	0.93	0.57	97	16
D. f. 6	1.84	0.86	0.67	0.87	0.57	99	19
M	1.98	0.90	0.71	0.93	0.60	96.67	17.17
SD	0.148	0.037	0.024	0.073	0.034	1.211	2.229
Min	1.83	0.86	0.67	0.87	0.57	96	14
Max	2.21	0.95	0.74	1.05	0.66	99	20

**Figure 5** Principal component analysis (PCA) on the shells of *Dalmatinella* based on the measurements given in Table 2.

nearly vestigial, female organs not different from the ones of *D. fluviatilis* (Radoman 1973, 1974, 1983–Falniowski & Szarowska 2013). Penis (Fig.

7) with two small, non-glandular lobes, lobes less prominent than in *D. fluviatilis* (Radoman 1973, 1983–Falniowski & Szarowska 2013).




**Figure 6** Renal and pallial section of the female reproductive organs of *Dalmatinella simonae*, a paratype (bc – bursa copulatrix, cbc – duct of bursa copulatrix, ga – albuminoid gland, gn – nidamental gland, gp – gonoporus, ov – oviduct, ovl – loop of (renal) oviduct, rs<sub>1</sub> rs<sub>2</sub> – receptaculum seminis, terminology after Radoman (1973): rs<sub>1</sub> – distal, rs<sub>2</sub> – proximal); bar equals 0.5mm.

*Derivation of name* The specific epithet *simonae* refers to Simona Prevorčnik from the University of Ljubljana, our friend and contributor to our study of stygobiont Truncatelloidea of the Balkans.

*Distribution and habitat* Found at type locality (Ploče, S part of lake Sladinac, Bačina lakes, Fig. 1 – site 3), but also in Cetina River (Donja Rošca and Trnbusi, Fig. 1 – sites 1 and 2), on stones of the banks of the river, dam reservoirs or lakes.

#### MOLECULAR DISTINCTNESS AND RELATIONSHIPS OF *DALMATINELLA SIMONAE*

We obtained six new sequences of COI (457 bp, GenBank Accession Numbers MT773271–MT773276). The tests by Xia *et al.* (2003) revealed no saturation. In all analyses, the topologies of the resulting phylograms were identical in




**Figure 7** Penis of *Dalmatinella simonae*, two paratypes (x – tip, lb – non-glandular lobe); bar equals 0.5mm.

both the ML and BI. The new sequences clearly belonged to the genus *Dalmatinella* (Fig. 8). The sequences of cytochrome oxidase COI differ by the p-distance 0.039 between *Dalmatinella simonae* and *D. fluviatilis*, confirming species distinctness of the two taxa. Within the species there was nearly no haplotype diversity, the p-distance between the haplotype from the locality 1 and the haplotype from the localities 2 and 3 (the same haplotype at the latter two localities) was only 0.003 (Fig. 8). *Dalmatinella* clearly belongs to the Sadlerianinae, but exact phylogenetic relationship with other genera is uncertain due to low bootstrap values.

#### DISCUSSION

Falniowski (1987) in his monograph of the Polish Hydrobioidea illustrated and described the variability of the characters used in the truncatelloid taxonomy (male and female reproductive organs, ctenidium, osphradium, soft part pigmentation, etc.) and considered such characters, like the proportions of the penis, receptacula, bursa, etc., much more variable than it had been usually noticed in the literature. Szarowska (2006) reconstructed character evolution in the Balkan Truncatelloidea, and found long-lasting



**Figure 8** Phylogenetic relationships of *Dalmatinella*: maximum likelihood (ML) phylogram based on COI; *Ecrobia maritima* (Hydrobiinae) given as outgroup.

evolutionary stability of the presence and topological position of bursae, receptacula and penis outgrowths, but high variability of their shape, proportions and relative size. There are several factors shaping this variability, coupled with relative simplicity of these structures (Szarowska & Falniowski 2008–Falniowski 2018), but slight differences observed between *Dalmatinella fluviatilis* and *D. simonae* are not surprising.

Our results confirm the opinion of Falniowski & Szarowska that *Dalmatinella* inhabiting Neretva River may not belong to *D. fluviatilis*. Our findings at two localities in Cetina River suggest more continuous geographic distribution of the genus.

Interestingly, the geographic distance between the most distant localities of *D. simonae* (1 and 3) is similar to the one between the westernmost locality of *D. simonae* (locality 1) and the locality inhabited by *D. fluviatilis*. At the same time, the genetic p-distance between the localities 1 and 2/3 approaches 0.003, while the p-distance between the species equals 0.039.

The above value of the genetic distance fall within the zone characteristic either of weakly marked interspecific values or high intraspecific variation, but rather characterise interspecific values (e.g. Perez *et al.*, 2005; Falniowski *et al.*, 2007; Szarowska *et al.*, 2007; Falniowski & Szarowska,



2011a; Szarowska & Falniowski, 2013). Bichain *et al.* (2007) reported the threshold value 0.015 in the west-European *Bythinella* species. In case of our *Dalmatinella* the virtual lack of intraspecific genetic variation between three localities of *D. simonae* supports the interspecies range of the observed p-distance 0.039. Obviously, the evolution of a gene need not necessarily reflect the evolution of species (e.g. Avise, 2000), but in the case of *Dalmatinella* the genetic differences is coupled with the differences in the proportions of the shell, clearly illustrated by PCA, and by different habitus of the penis, which confirm species distinctness of the two taxa.

#### ACKNOWLEDGMENTS

The study was supported by a grant from the National Science Centre 2017/25/B/NZ8/01372 to Andrzej Falniowski.

#### REFERENCES

- AVISE JC 2000 Phylogeography. *The History and Formation of Species*. Cambridge; London: Harvard University Press.
- BERAN L, HOFMAN S & FALNIOWSKI A 2015 *Tanousia zrmanjiae* (Brusina, 1866) (Caenogastropoda: Truncatelloidea: Hydrobidae): A living fossil *Folia Malacologica* **23**: 263–271.
- BERAN L, OSIKOWSKI A, HOFMAN S & FALNIOWSKI A 2016 *Islamia zermanica* (Radoman, 1973) (Caenogastropoda: Hydrobidae): morphological and molecular distinctness *Folia Malacologica* **24**: 25–30.
- BICHAIN J-M, GAUBERT P, SAMADI S & BOISSELIER-DUBAYLE M-C 2007 A gleam in the dark: Phylogenetic species delimitation in the confusing spring-snail genus *Bythinella* Moquin-Tandon, 1856 (Gastropoda: Risssooidea: Amnicolidae) *Molecular Phylogenetics and Evolution* **45**: 927–941.
- EDGAR RC 2004 MUSCLE: multiple sequence alignment with high accuracy and high throughput *Nucleic Acids Research* **32**: 1792–1797.
- FALNIOWSKI A 1987 Hydrobioidea of Poland (Prosobranchia: Gastropoda) *Folia Malacologica* **1**: 1–122.
- FALNIOWSKI A 2003 Metody numeryczne w taksonomii [Numerical techniques in taxonomy]. Kraków: Wydawnictwo Uniwersytetu Jagiellońskiego.
- FALNIOWSKI A 2018 Species Distinction and Speciation in Hydrobioid Gastropods (Mollusca: Caenogastropoda: Truncatelloidea) *Archives of Zoological Studies* **1**: 003.
- FALNIOWSKI A & SZAROWSKA M 2011a Radiation and phylogeography in a spring snail *Bythinella* (Mollusca: Gastropoda: Risssooidea) in continental Greece *Annales Zoologici Fennici* **48**: 67–90.
- FALNIOWSKI A & SZAROWSKA M 2011b A new genus and new species of valvatiform hydrobiid (Risssooidea; Caenogastropoda) from Greece *Molluscan Research* **31**: 189–199.
- FALNIOWSKI A & SZAROWSKA M 2013 Phylogenetic relationships of *Dalmatinella fluviatilis* Radoman, 1973 (Caenogastropoda: Risssooidea) *Folia Malacologica* **21**: 1–7.
- FALNIOWSKI A, SZAROWSKA M & GRZMIL P 2007 *Daphniola* Radoman, 1973 (Gastropoda: Hydrobiidae): shell biometry, mtDNA, and the Pliocene flooding *Journal of Natural History* **41**: 2301–2311.
- FALNIOWSKI A, SZAROWSKA M, GLÖER P & PESIC V 2012 Molecules vs morphology in the taxonomy of the *Radomaniola/Grossuana* group of Balkan Risssooidea (Mollusca: Caenogastropoda) *Journal of Conchology* **41**: 19–36.
- FALNIOWSKI A, PESIC V & GLÖER P 2014 *Montenegrospeum* Pešić et Glöer, 2013: a representative of Moitessieriidae? *Folia Malacologica* **22**: 263–268.
- FALNIOWSKI A, GEORGIEV D, OSIKOWSKI A & HOFMAN S 2016 Radiation of *Grossuana* Radoman, 1973 (Caenogastropoda: Truncatelloidea) in the Balkans *Journal of Molluscan Studies* **82**: 305–313.
- HALL TA 1999 BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT *Nucleic Acids Symposium Series* **41**: 95–98.
- HERSHLER R & PONDER WF 1998 A review of morphological characters of hydrobioid snails *Smithsonian Contribution to Zoology* **600**: 1–55.
- METSALU T & JAAK V 2015 Clustvis: a web tool for visualizing clustering of multivariate data using Principal Component Analysis and heatmap. *Nucleic Acids Research* **43**: 566–570.
- MILLER MA, PFEIFFER W & SCHWARTZ T 2010 Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In: Proceedings of the Gateway Computing Environments Workshop (GCE), 14 Nov, New Orleans, LA: 1–8.
- NYLANDER JAA 2004 MrModeltest v.2. Program distributed by the author. Uppsala: Evolutionary Biology Centre, Uppsala University.
- OSIKOWSKI A, HOFMAN S, GEORGIEV D, KALCHEVA S & FALNIOWSKI A 2016 Aquatic snails *Ecrobia maritima* (Milaschewitsch, 1916) and *E. ventrosa* (Montagu, 1803) (Caenogastropoda: Hydrobiidae) in the east Mediterranean and Black Sea *Annales Zoologici* **66**: 477–486.
- OSIKOWSKI A, HOFMAN S, RYSIEWSKA A, SKET B, PREVORCNIK S & FALNIOWSKI A 2018 A case of biodiversity overestimation in the Balkan *Belgrandiella* A. J. Wagner, 1927 (Caenogastropoda: Hydrobiidae): molecular divergence not paralleled by high morphological variation *Journal of Natural History* **52**: 323–344.
- PEREZ KE, PONDER WF, COLGAN DJ, CLARK SA & LYDEARD C 2005 Molecular phylogeny and biogeography of spring-associated hydrobiid snails of the Great Artesian Basin, Australia *Molecular Phylogenetics and Evolution* **34**: 545–556.

- RADOMAN P 1973 New classification of fresh and brackish water Prosobranchia from the Balkans and Asia Minor *Posebna Izdanja, Prirodnjacki Musej u Beogradu* **32**: 1–30.
- RADOMAN P 1974 *Dalmaninella* and *Vinodolia*. New fresh water genera of Prosobranchia from the Yugoslav seaside *Glasnik Prirodnjackog Museja, Beograd series B* **29**: 41–44.
- RADOMAN P 1983 Hydrobioidea a super family of Prosobranchia (Gastropoda). I. Systematics. *Serbian Academy of Sciences and Arts, Monographs* **547**, Department of Sciences **57**: 1–256.
- RAMBAUT A & DRUMMOND AJ 2009 Tracer v1.5. <<http://beast.bio.ed.ac.uk/Tracer>>.
- RONQUIST F, TESLENKO M, VAN DER MARK P, AYRES DL, DARLING A, HÖHNA S, LARGET B, LIU L, SUCHARD MA & HUELSENBECK JP 2012 Efficient Bayesian phylogenetic inference and model choice across a large model space *Systematic Biology* **61**: 539–542.
- ROHLF FJ 1998 *NTSYSpc, numerical taxonomy and multivariate analysis system. Version 2.0*. [Computer software and manual] Exeter Software, Seatuket, NY.
- RUEDEN CT, SCHINDELIN J, HINER MC, DEZONIA BE, WALTER AE, ARENA ET & ELICEIRI KW 2017 ImageJ2: ImageJ for the next generation of scientific image data *BMC Bioinformatics* **18**: 529.
- RSYIEWSKA A, GEORGIEV D, OSIKOWSKI A, HOFMAN S & FALNIOWSKI A 2016 *Pontobelgrandiella* Radoman, 1973 (Caenogastropoda: Hydrobiidae): A recent invader of subterranean waters? *Journal of Conchology* **42**: 193–203.
- RSYIEWSKA A, PREVORCNIK S, OSIKOWSKI A, HOFMAN S, BERAN L & FALNIOWSKI A 2017 Phylogenetic relationships in *Kerkia* and introgression between *Hauffenia* and *Kerkia* (Caenogastropoda: Hydrobiidae) *Journal of Zoological Systematics and Evolutionary Research* **55**: 106–117.
- STAMATAKIS A 2014 RAxML Version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies *Bioinformatics* **30**: 1312–1313.
- SZAROWSKA M 2006 Molecular phylogeny, systematics and morphological character evolution in the Balkan Rissooidea (Caenogastropoda) *Folia Malacologica* **14**: 99–168.
- SZAROWSKA M & FALNIOWSKI A 2008 *There is no philosopher's stone: coup de grace for the morphology-based systematics in the rissooidean gastropods?* 5<sup>th</sup> Congress of the European Malacological Societies, Ponta Delgada: 28.
- SZAROWSKA M & FALNIOWSKI A 2011 An unusual, flagellum-bearing hydrobiid snail (Gastropoda: Rissooidea: Hydrobiidae) from Greece, with descriptions of a new genus and a new species *Journal of Natural History* **45**: 2231–2246
- SZAROWSKA M & FALNIOWSKI A 2013 Species distinctness of *Sadleriana robici* (Clessin, 1890) (Gastropoda: Rissooidea) *Folia Malacologica* **21**: 127–133.
- SZAROWSKA M & FALNIOWSKI A 2014 *Horatia Bourguignat, 1887*: is this genus really phylogenetically very close to *Radomaniola* Szarowska, 2006 (Caenogastropoda: Truncatelloidea)? *Folia Malacologica* **22**: 31–39.
- SZAROWSKA M, GRZMIL P, FALNIOWSKI A & SIRBU I 2007 *Grossuana codreanui* (Grossu, 1946) and the phylogenetic relationships of the East Balkan genus *Grossuana* (Radoman, 1973) (Gastropoda: Rissooidea) *Hydrobiologia* **579**: 379–391.
- SZAROWSKA M, HOFMAN S, OSIKOWSKI A & FALNIOWSKI A 2014a *Daphniola* Radoman, 1973 (Caenogastropoda: Truncatelloidea) at east Aegean islands *Folia Malacologica* **22**: 269–275.
- SZAROWSKA M, HOFMAN S, OSIKOWSKI A & FALNIOWSKI A 2014b Divergence preceding Island formation among Aegean insular populations of the freshwater snail genus *Pseudorientalia* (Caenogastropoda: Truncatelloidea) *Zoological Science* **31**: 680–686.
- SZAROWSKA M, OSIKOWSKI A, HOFMAN S & FALNIOWSKI A 2016 *Pseudamnicola* Paulucci, 1878 (Caenogastropoda: Truncatelloidea) from the Aegean Islands: a long or short story? *Organism Diversity and Evolution* **16**: 121–139.
- TAMURA K, STECHER G, PETERSON D, FILIPSKI A & KUMAR S 2013 MEGA6: molecular evolutionary genetics analysis version 6.0 *Molecular Biology and Evolution* **30**: 2725–2729.
- WILKE T, DAVIS GM, FALNIOWSKI A, GIUSTI F, BODON M & SZAROWSKA M 2001 Molecular systematics of Hydrobiidae (Mollusca: Gastropoda: Rissooidea): testing monophyly and phylogenetic relationships *Proceedings of the Academy of Natural Sciences of Philadelphia* **151**: 1–21.
- XIA X 2000 *Data analysis in molecular biology and evolution* Kluwer Academic Publishers, Boston, Dordrecht & London.
- XIA X 2013 DAMBE: A comprehensive software package for data analysis in molecular biology and evolution *Molecular Biology and Evolution* **30**: 1720–1728.
- XIA X, XIE Z, SALEMI M, CHEN L & WANG Y 2003 An index of substitution saturation and its application *Molecular Phylogenetics and Evolution* **26**: 1–7.